

Journées

ÉQUATIONS AUX DÉRIVÉES PARTIELLES

Biarritz, 6 juin–10 juin 2011

Benjamin Dodson

Global well-posedness and scattering for the mass-critical NLS

J. É. D. P. (2011), Exposé n° IV, 11 p.

<http://jedp.cedram.org/item?id=JEDP_2011____A4_0>

cedram

*Article mis en ligne dans le cadre du
Centre de diffusion des revues académiques de mathématiques
<http://www.cedram.org/>*

Global well-posedness and scattering for the mass-critical NLS

Benjamin Dodson

Suppose $u(t, x)$ is a solution to the nonlinear partial differential equation

$$\begin{aligned} iu_t + \Delta u &= \mu |u|^{4/d} u, \\ u(0, x) &= u_0 \in L^2(\mathbf{R}^d), \end{aligned} \tag{0.1}$$

$\mu = \pm 1$, $\mu = +1$ refers to the defocusing case and $\mu = -1$ refers to the focusing case.

Definition 0.1. (0.1) is said to be globally well - posed if a solution $u(t, x)$ to (0.1) exists for all time,

$$u(t, x) \in C_t^0(\mathbf{R}; L^2(\mathbf{R}^d)) \cap L_{t,loc}^{\frac{2(d+2)}{d}}(\mathbf{R}; L^{\frac{2(d+2)}{d}}(\mathbf{R}^d)), \tag{0.2}$$

and a solution to (0.1) depends continuously on u_0 in the $L^2(\mathbf{R}^d)$ topology.

Definition 0.2. A global solution to (0.1) is said to scatter if there exist $u_{\pm} \in L^2(\mathbf{R}^d)$ such that

$$\|u(t, x) - e^{it\Delta} u_+\|_{L^2(\mathbf{R}^d)} \rightarrow 0, \tag{0.3}$$

as $t \rightarrow +\infty$ and

$$\|u(t, x) - e^{it\Delta} u_-\|_{L^2(\mathbf{R}^d)} \rightarrow 0 \tag{0.4}$$

as $t \rightarrow -\infty$. Additionally we say a solution to (0.1) scatters forward in time if it satisfies (0.3) and backward in time if it satisfies (0.4).

The first progress toward proving well - posedness of (0.1) was

Theorem 0.1. (0.1) is locally well - posed on $[-T, T]$ for some $T(\|u_0\|_{H^1(\mathbf{R}^d)}) > 0$.

Proof: See [6]. \square

Furthermore, it is possible to use conserved quantities of (0.1) to upgrade theorem 0.1 to global well - posedness. (0.1) has the conserved quantities mass,

$$M(u(t)) = \int |u(t, x)|^2 dx = M(u(0)), \tag{0.5}$$

and energy

$$E(u(t)) = \frac{1}{2} \int |\nabla u(t, x)|^2 dx + \frac{\mu d}{2(d+2)} \int |u(t, x)|^{\frac{2(d+2)}{d}} dx. \quad (0.6)$$

In the defocusing case (0.6) is positively definite, which implies $\|u(t)\|_{H^1(\mathbf{R}^d)}$ is uniformly bounded by $E(u(0))$ which is finite by the Sobolev embedding theorem. By (0.1) (0.1) is globally well - posed for $u_0 \in H^1(\mathbf{R}^d)$, $\mu = +1$.

In the focusing case (0.6) is not positive definite. Therefore having $E(u(0))$ finite is not enough to prove global well - posedness because $\|u(t)\|_{H^1(\mathbf{R}^d)}$ and $\|u(t)\|_{L^{\frac{2(d+2)}{d}}(\mathbf{R}^d)}$ can and do blow up at the same rate, precisely canceling to maintain conservation of energy.

For $\|u(t)\|_{L^2(\mathbf{R}^d)}$ below a certain threshold it is still possible to prove global well - posedness and scattering in the case when $\mu = -1$ using the Gagliardo - Nirenberg inequality.

Theorem 0.2. *If Q is the positive solution to the elliptic partial differential equation*

$$\Delta Q + Q^{1+4/d} = Q, \quad (0.7)$$

the Sobolev embedding theorem has the best constant

$$\|u\|_{L_x^{\frac{2(d+2)}{d}}(\mathbf{R}^d)}^{\frac{2(d+2)}{d}} \leq \frac{\|u\|_{L^2(\mathbf{R}^d)}^{4/d}}{\|Q\|_{L^2(\mathbf{R}^d)}^{4/d}} \|\nabla u\|_{L^2(\mathbf{R}^d)}^2. \quad (0.8)$$

Proof: See [30], [43], [44], and [5]. \square

Combining theorem 0.2 with (0.5) proves (0.1) when $\|u_0\|_{L^2(\mathbf{R}^d)} < \|Q\|_{L^2(\mathbf{R}^d)}$. Furthermore, by (0.7)

$$u(t, x) = e^{it}Q(x) \quad (0.9)$$

is a solution to (0.1) when $\mu = -1$. This is a solution that certainly fails to scatter. Applying the conformal symmetry

Theorem 0.3. *u is a solution to (0.1) if and only if*

$$v(t, x) = \frac{1}{|t|^{d/2}} u\left(-\frac{1}{t}, \frac{x}{t}\right) e^{i|x|^2/t} \quad (0.10)$$

solves (0.1).

We obtain a solution to (0.1) that fails to be globally well - posed.

Furthermore, consider the variance

$$\int |x|^2 |u(t, x)|^2 dx. \quad (0.11)$$

$$\frac{d^2}{dt^2} \int |x|^2 |u(t, x)|^2 dx = 16E(u(t)). \quad (0.12)$$

If $u(0) \in H^1(\mathbf{R}^d)$, $0 > E(u(t)) > -\infty$, and $\int |x|^2 |u(0, x)|^2 dx < \infty$, then the variance is concave down in time, which implies that $\int |x|^2 |u(t, x)|^2 dx$ will cross the real axis

twice. Since (0.11) is positive definite, this implies a solution to (0.1) can only exist in both directions for finite time. Such solutions are relatively straightforward to construct when $\|u_0\|_{L^2(\mathbf{R}^d)} > \|Q\|_{L^2(\mathbf{R}^d)}$.

The local well - posedness result in theorem 0.1 was substantially improved to

Theorem 0.4. (0.1) is locally well - posed on $[-T, T]$ for $u_0 \in L^2(\mathbf{R}^d)$, $T(u_0) > 0$, where T depends on the profile of u_0 , not just its size.

Proof: See [6] and [7]. \square

In this paper we sketch the proof of the natural extension of theorem 0.4,

Theorem 0.5. (0.1) is globally well - posed and scattering for all $u_0 \in L^2(\mathbf{R}^d)$, $\mu = +1$. (0.1) is globally well - posed and scattering for $\mu = -1$, $\|u_0\|_{L^2(\mathbf{R}^d)} < \|Q\|_{L^2(\mathbf{R}^d)}$.

Previous Results:

Theorem 0.6. (0.1) is globally well - posed and scattering for $\mu = +1$, $u_0 \in L^2(\mathbf{R}^d)$ and for $\mu = -1$, $\|u_0\|_{L^2(\mathbf{R}^d)} < \|Q\|_{L^2(\mathbf{R}^d)}$ when u_0 is radial, $d = 2$.

Proof: See [24]. \square

Theorem 0.7. (0.1) is globally well - posed and scattering for $\mu = +1$, $u_0 \in L^2(\mathbf{R}^d)$ when u_0 is radial, $d \geq 3$.

Proof: See [37]. \square

Theorem 0.8. (0.1) is globally well - posed and scattering for $\mu = -1$, $\|u_0\|_{L^2(\mathbf{R}^d)} < \|Q\|_{L^2(\mathbf{R}^d)}$ when u_0 is radial, $d \geq 3$.

Proof: See [26]. \square

Conjecture: If (0.1) is not globally well - posed and scattering, and $\|u_0\|_{L^2(\mathbf{R}^d)} = \|Q\|_{L^2(\mathbf{R}^d)}$, then

$$u(t, x) = G \cdot e^{it}Q(x) \quad (0.13)$$

where $G = S^1 \times (0, \infty) \times \mathbf{R}^d \times \mathbf{R}^d$ is a group of symmetries acting on solutions to (0.1), or a conformal symmetry of (0.13).

G is generated by four symmetries which act on solutions of (0.1), multiplication,

$$u(t, x) \mapsto e^{i\theta}u(t, x), \quad (0.14)$$

for $\theta \in \mathbf{R}$, scaling,

$$u(t, x) \mapsto \frac{1}{\lambda^{d/2}}u\left(\frac{t}{\lambda^2}, \frac{x}{\lambda}\right), \quad (0.15)$$

translation,

$$u(t, x) \mapsto u(t, x - x_0), \quad (0.16)$$

and Galilean invariance

$$u(t, x) \mapsto e^{-it|\xi_0|^2}e^{ix \cdot \xi_0}u(t, x - 2t\xi_0). \quad (0.17)$$

Let

$$A_\mu(m) = \sup\{\|u\|_{L_{t,x}^{\frac{2(d+2)}{d}}(\mathbf{R} \times \mathbf{R}^d)} : \|u(t)\|_{L_x^2(\mathbf{R}^d)} = m, u \text{ solves (0.1)}\}. \quad (0.18)$$

To prove theorem 0.1 it suffices to prove $A_\mu(m) < \infty$ for all m in the defocusing case and for $m < \|Q\|_{L^2(\mathbf{R}^d)}$ in the focusing case.

Theorem 0.9. $A_\mu(m)$ is a continuous function of m .

Proof: See [35].

This already implies a small data result for $m < \epsilon(d, \mu)$ because $A_\mu(0) = 0$. Moreover,

$$\{m : A_\mu(m) = \infty\} \quad (0.19)$$

is a closed set so if (0.19) is nonempty then it possesses a least element m_0 .

Theorem 0.10. Suppose

$$\|u_n(t)\|_{L_x^2(\mathbf{R}^d)} \nearrow m_0 \quad (0.20)$$

$$\|u_n\|_{L_{t,x}^{\frac{2(d+2)}{d}}(t \geq 0)} \nearrow \infty, \quad \|u_n\|_{L_{t,x}^{\frac{2(d+2)}{d}}(t \leq 0)} \nearrow \infty. \quad (0.21)$$

Then $u_n(t)$ has a subsequence that converges to $u(t)$ in $L^2(\mathbf{R}^d)/G$, $u(t) : I \subset \mathbf{R} \rightarrow \mathbf{C}$ is a solution to (0.1), I an open set.

$$\|u(t)\|_{L_{t,x}^{\frac{2(d+2)}{d}}(t \geq 0)} = \|u(t)\|_{L_{t,x}^{\frac{2(d+2)}{d}}(t \leq 0)} = \infty. \quad (0.22)$$

Moreover, $\{u(t) : t \in I\}$ lies in a compact subset of $L^2(\mathbf{R}^d)/G$. By the Arzela-Ascoli theorem there exist

$$x(t), \xi(t) : I \rightarrow \mathbf{R}^d, \quad (0.23)$$

$$N(t) : I \rightarrow (0, \infty), \quad (0.24)$$

such that for all $\eta > 0$ there exists $C(\eta) < \infty$ such that

$$\int_{|x-x(t)| \geq \frac{C(\eta)}{N(t)}} |u(t, x)|^2 dx + \int_{|\xi-\xi(t)| \geq C(\eta)N(t)} |\hat{u}(t, \xi)|^2 d\xi < \eta. \quad (0.25)$$

Proof: See [36]. \square

Because $u(t)$ lies in a precompact set we can take a limit of $u(t_n)$, $t_n \in I$, in $L^2(\mathbf{R}^d)/G$ and obtain an even more special solution to (0.1).

Theorem 0.11. If theorem 0.1 fails then there exists a solution to (0.1) satisfying $N(0) = 1$, $u(t)$ exists on $[0, \infty)$, $N(t) \leq 1$ on $[0, \infty)$, $x(0) = \xi(0) = 0$,

$$|\xi'(t)|, |N'(t)| \lesssim_{m_0, d} N(t)^3, \quad (0.26)$$

$$\|u\|_{L_{t,x}^{\frac{2(d+2)}{d}}([0, \infty) \times \mathbf{R}^d)} = \|u\|_{L_{t,x}^{\frac{2(d+2)}{d}}((\inf(I), 0] \times \mathbf{R}^d)} = \infty. \quad (0.27)$$

Proof: See [25]. \square

We consider two cases separately,

$$\int_0^\infty N(t)^3 dt < \infty, \quad (0.28)$$

and

$$\int_0^\infty N(t)^3 dt = \infty. \quad (0.29)$$

Theorem 0.12. *If $\int_0^\infty N(t)^3 dt = K$, then for all $0 \leq s < 1 + \frac{4}{d}$,*

$$\|u\|_{L_t^\infty \dot{H}_x^s([0, \infty) \times \mathbf{R}^d)} \lesssim_{m_0, d} K^s. \quad (0.30)$$

This is enough to exclude (0.28) in the defocusing case and in the focusing case for mass below the mass of the ground state. In both cases $E(u(0)) \geq \delta > 0$. (0.26) and (0.28) imply $N(t) \rightarrow 0$ as $t \rightarrow \infty$. So when $\xi(t) \equiv 0$ it is easy to see $N(t) \rightarrow 0$, Sobolev embedding, and (0.30) imply

$$E(u(t)) \rightarrow 0, \quad (0.31)$$

which contradicts conservation of energy. In the general case when $\xi(t)$ is free to move around, (0.26) implies that $|\xi(t)| \lesssim_{m_0, d} K$ for all $t \in [0, \infty)$. For T sufficiently large $N(T)$ is very small. After making a Galilean transformation sending $\xi(T)$ to 0, this implies $E(u(T))$ is very small. Because $|\xi(T)| \lesssim_{m_0, d} K$ this transformation preserves (0.30). On the other hand, because L^p is Galilean invariant, after any Galilean transformation

$$E(u(0)) \geq \delta > 0. \quad (0.32)$$

This again contradicts conservation of energy.

Having completely ruled out case (0.28) we turn to case (0.29) when $\mu = +1$. We use the interaction Morawetz estimate

Theorem 0.13. *If $\mu = +1$,*

$$\| |\nabla|^{\frac{3-d}{2}} |u(t, x)|^2 \|_{L_{t,x}^2([0, T] \times \mathbf{R}^d)}^2 \lesssim_{m_0, d} \int_0^T \partial_t M(t) dt, \quad (0.33)$$

where

$$M(t) = \int \frac{(x-y)_j}{|x-y|} \text{Im}[\bar{u}(t, x) \partial_j u(t, x)] |u(t, y)|^2 dx dy. \quad (0.34)$$

Because the solution $u(t)$ need not possess any additional regularity, we truncate in frequency.

Theorem 0.14. *Suppose $\int_0^T N(t)^3 dt = K$, choose C sufficiently large so that*

$$\int_0^T |\xi'(t)| dt \ll CK, \quad (0.35)$$

which is always possible by (0.26). Let

$$M(t) = \int \frac{(x-y)_j}{|x-y|} \text{Im}[P_{\leq CK} \bar{u}(t,x) \partial_j P_{\leq CK} u(t,x)] |P_{\leq CK} u(t,y)|^2 dx dy. \quad (0.36)$$

Then

$$\int_0^T N(t)^3 dt \lesssim_{m_0,d} \int_0^T \partial_t M(t) dt, \quad (0.37)$$

and since $N(t) \leq 1$ for $t \in [0, \infty)$,

$$|M(t)| \lesssim_{m_0,d} o(K). \quad (0.38)$$

This rules out (0.29) in the case when $\mu = +1$ because K can be made arbitrarily large by taking T sufficiently large, giving the contradiction

$$K \lesssim o(K). \quad (0.39)$$

We now give a brief discussion of the proof of theorem 0.14. It is perhaps easiest to see that when $d \geq 4$, if u is a minimal mass blowup solution to (0.1),

$$N(t)^3 \lesssim_{m_0,d} \| |\nabla|^{\frac{3-d}{2}} |u(t,x)|^2 \|_{L_x^2(\mathbf{R}^d)}^2. \quad (0.40)$$

Indeed, when $d \geq 4$,

$$\| |\nabla|^{\frac{3-d}{2}} |u(t,x)|^2 \|_{L_x^2(\mathbf{R}^d)}^2 \sim_{m_0,d} \int \frac{1}{|x-y|^3} |u(t,x)|^2 |u(t,y)|^2 dx dy. \quad (0.41)$$

The spatial concentration in (0.25) implies (0.40). Because most of the mass is contained in $P_{\leq CK}$, we also have

$$\int_0^T N(t)^3 dt \lesssim_{m_0,d} \| |\nabla|^{\frac{3-d}{2}} |P_{\leq CK} u(t,x)|^2 \|_{L_{t,x}^2([0,T] \times \mathbf{R}^d)}^2. \quad (0.42)$$

(0.38) follows from (0.25), $N(t) \leq 1$, and the fact that the interaction Morawetz estimates are Galilean invariant.

$$i\partial_t(P_{\leq CK} u) + \Delta(P_{\leq CK} u) = \mu |P_{\leq CK} u|^{4/d} (P_{\leq CK} u) + \mu [P_{\leq CK} (|u|^{4/d} u) - |P_{\leq CK} u|^{4/d} (P_{\leq CK} u)]. \quad (0.43)$$

If we were able to drop

$$\mu [P_{\leq CK} (|u|^{4/d} u) - |P_{\leq CK} u|^{4/d} (P_{\leq CK} u)] \quad (0.44)$$

then the proof of (0.42) when $\mu = +1$ would be identical to the proof of (0.33). Therefore, most of the work in proving theorem 0.14 lies in showing that the error arising from (0.44) is bounded by $o(K)$. In fact, the error estimates are quite robust.

Theorem 0.15. *We can perform the same error estimates with $\frac{(x-y)_j}{|x-y|}$ replaced with $a(t, x-y)_j$ as long as*

$$|a(t, x)| \lesssim_{m_0,d} 1, \quad (0.45)$$

when $d = 2$,

$$|\nabla a(t, x)| \lesssim_{m_0,d} \frac{1}{|x|}, \quad (0.46)$$

and when $d = 1$,

$$\|\nabla a(t, x)\|_{L^1_x(\mathbf{R})} \lesssim_{m_0, d} 1, \quad (0.47)$$

$$a(t, x) = -a(t, -x), \quad (0.48)$$

and when $d = 2$,

$$\|\partial_t a(t, x)\|_{L^1(\mathbf{R}^2)} \lesssim_{m_0, d} 1. \quad (0.49)$$

Therefore it remains to construct an interaction Morawetz potential bounded below by $N(t)^3$ and which satisfies (0.45) - (0.49). We do this only in the case when $d = 1$ and u is radial. Suppose $\psi \in C^\infty(\mathbf{R})$, $\psi = \phi' \geq 0$, and

$$\begin{aligned} &= x, \quad |x| \leq 1, \\ \psi(x) &= \frac{3}{2}, \quad x > 2, \\ &= -\frac{3}{2}, \quad x < -2. \end{aligned} \quad (0.50)$$

Then let

$$M(t) = R \int \psi\left(\frac{x\tilde{N}(t)}{R}\right) \text{Im}[\bar{u}\partial_x u](t, x) dx, \quad (0.51)$$

such that for some $\delta > 0$

$$\delta N(t) \leq \tilde{N}(t) \leq N(t), \quad (0.52)$$

and

$$\int_0^T |\tilde{N}'(t)| dt \leq \delta_1 \int_0^T \tilde{N}(t) N(t)^2 dt. \quad (0.53)$$

Then

$$\partial_t M(t) = \tilde{N}(t) \int \phi\left(\frac{x\tilde{N}(t)}{R}\right) \left[\frac{1}{2} |\nabla u(t, x)|^2 - \frac{1}{6} |u(t, x)|^6 \right] dx \quad (0.54)$$

$$+ C \frac{\tilde{N}(t)^3}{R^2} \int \phi''\left(\frac{x\tilde{N}(t)}{R}\right) |u(t, x)|^2 dx \quad (0.55)$$

$$+ R \int \phi\left(\frac{x\tilde{N}(t)}{R}\right) x \tilde{N}'(t) \text{Im}[\bar{u}\partial_x u](t, x) dx. \quad (0.56)$$

We choose $\tilde{N}(t)$ to be a sufficiently slowly varying (0.53) envelope for $N(t)$ which allows us to absorb (0.55) into (0.54) for $R(m_0)$ sufficiently large and for $\delta_1(R)$ sufficiently small we can absorb (0.56) into (0.54). This completes the proof of the focusing case.

References

- [1] J. Bourgain “Fourier transform restriction phenomena for certain lattice subsets and applications to nonlinear evolution equations. I. Schrödinger equations” *Geom. Funct. Anal.* **3** (1993): 2, 107 – 156.
- [2] J. Bourgain “Fourier transform restriction phenomena for certain lattice subsets and applications to nonlinear evolution equations. II. The KdV-equation” *Geom. Funct. Anal.* **3** (1993): 3, 209–262.
- [3] J. Bourgain. “Refinements of Strichartz’ inequality and applications to 2D-NLS with critical nonlinearity.” *International Mathematical Research Notices*, **5** (1998):253 – 283.
- [4] J. Bourgain. “Global Solutions of Nonlinear Schrödinger Equations” *American Mathematical Society Colloquium Publications*, 1999.
- [5] H. Berestycki and P.L. Lions, two authors *Existence d’ondes solitaires dans des problèmes nonlinéaires du type Klein-Gordon*, Comptes Rendus Hebdomadaires des Séances de l’Académie des Sciences. Séries A et B, **288** no. 7 (1979), A395 - A398.
- [6] T. Cazenave and F. B. Weissler, *The Cauchy problem for the nonlinear Schrödinger equation in H^1* , Manuscripta Math., **61** (1988), 477–494.
- [7] T. Cazenave and F. B. Weissler, two authors "The Cauchy problem for the critical nonlinear Schrödinger equation in H^s ", *Nonlinear Anal.*, **14** (1990), 807–836.
- [8] J. Colliander, M. Grillakis, and N. Tzirakis. "Improved interaction Morawetz inequalities for the cubic nonlinear Schrödinger equation on \mathbf{R}^2 ." *International Mathematics Research Notices. IMRN*, **23** (2007): 90 - 119.
- [9] J. Colliander, M. Grillakis, and N. Tzirakis. "Tensor products and correlation estimates with applications to nonlinear Schrödinger equations" *Communications on Pure and Applied Mathematics*, **62** no. 7 (2009) : 920 - 968
- [10] J. Colliander, M. Keel, G. Staffilani, H. Takaoka, and T. Tao. "Almost conservation laws and global rough solutions to a nonlinear Schrödinger equation." *Mathematical Research Letters*, **9** (2002):659 – 682.
- [11] J. Colliander, M. Keel, G. Staffilani, H. Takaoka, and T. Tao. "Global existence and scattering for rough solutions of a nonlinear Schrödinger equation on \mathbf{R}^3 " *Communications on pure and applied mathematics*, **21** (2004) : 987 - 1014
- [12] J. Colliander, M. Keel, G. Staffilani, H. Takaoka, and T. Tao. "Resonant decompositions and the I-method for cubic nonlinear Schrödinger equation on \mathbf{R}^2 ." *Discrete and Continuous Dynamical Systems A*, **21** (2007):665 – 686.
- [13] J. Colliander, M. Keel, G. Staffilani, H. Takaoka, and T. Tao. "Global existence and scattering for the energy - critical nonlinear Schrödinger equation on \mathbf{R}^3 " *Annals of Mathematics. Second Series*, **167** (2008) : 767 - 865

- [14] J. Colliander and T. Roy, *Bootstrapped Morawetz Estimates and Resonant Decomposition for Low Regularity Global solutions of Cubic NLS on \mathbf{R}^2* , preprint, *arXiv:0811.1803*,
- [15] B. Dodson, *Global well-posedness and scattering for the defocusing L^2 -critical nonlinear Schrödinger equation when $d \geq 3$* , preprint, *arXiv:0912.2467v1*,
- [16] B. Dodson, *Global well-posedness and scattering for the defocusing L^2 -critical nonlinear Schrödinger equation when $d = 1$* , preprint, *arXiv:1010.0040v2*,
- [17] B. Dodson, *Global well-posedness and scattering for the defocusing L^2 -critical nonlinear Schrödinger equation when $d = 2$* , preprint, *arXiv:1006.1375v2*,
- [18] B. Dodson, *Global well-posedness and scattering for the mass critical nonlinear Schrödinger equation with mass below the mass of the ground state*, preprint, *arXiv:1104.1114v2*,
- [19] P. Germain, N. Masmoudi, and J. Shatah, *Global solutions for 2D quadratic Schrödinger equations*, preprint, *arXiv:1001.5158v1*,
- [20] M. Hadac and S. Herr and H. Koch “Well-posedness and scattering for the KP-II equation in a critical space” *Ann. Inst. H. Poincaré Anal. Non Linéaire* **26** (2009): 3, 917–941.
- [21] C. Kenig and F. Merle “Global well-posedness, scattering, and blow-up for the energy-critical, focusing nonlinear Schrödinger equation in the radial case,” *Inventiones Mathematicae* **166** (2006): 3, 645–675.
- [22] C. Kenig and F. Merle “Scattering for $\dot{H}^{1/2}$ bounded solutions to the cubic, defocusing NLS in 3 dimensions,” *Transactions of the American Mathematical Society* **362** (2010): 4, 1937 – 1962.
- [23] M. Keel and T. Tao “Endpoint Strichartz Estimates” *American Journal of Mathematics* **120** (1998): 4 - 6, 945 – 957.
- [24] R. Killip, T. Tao, and M. Visan “The cubic nonlinear Schrödinger equation in two dimensions with radial data” *Journal of the European Mathematical Society*, to appear.
- [25] R. Killip and M. Visan “Nonlinear Schrödinger Equations at Critical Regularity” *Unpublished lecture notes*, Clay Lecture Notes (2009): <http://www.math.ucla.edu/visan/lecturenotes.html>.
- [26] R. Killip, M. Visan, and X. Zhang “The mass-critical nonlinear Schrödinger equation with radial data in dimensions three and higher” *Annals in PDE*, textbf1, no. 2 (2008) 229 - 266
- [27] H. Koch and D. Tataru “Dispersive estimates for principally normal pseudo-differential operators” *Communications on Pure and Applied Mathematics* **58** no. 2 (2005): 217 - 284

- [28] H. Koch and D. Tataru “A priori bounds for the 1D cubic NLS in negative Sobolev spaces” *Int. Math. Res. Not. IMRN* **16** (2007): Art. ID rnm053, 36.
- [29] H. Koch and D. Tataru, *Energy and local energy bounds for the 1-D cubic NLS equation in $H^{-1/4}$* , preprint, *arXiv:1012.0148*,
- [30] M. K. Kwong, *Uniqueness of positive solutions of $\Delta u - u + u^p = 0$ in \mathbf{R}^n* , *Archive for Rational Mechanics and Analysis* **105** no. 3 (1989), 243 - 266.
- [31] T. Ozawa and Y. Tsutsumi, *Space-time estimates for null gauge forms and nonlinear Schrödinger equations*, *Differential Integral Equations*, **11** no. 2 (1998), 201–222.
- [32] F. Planchon and L. Vega “Bilinear virial identities and applications” *Annales Scientifiques de l’École Normale Supérieure* **42**, no. 2 (2009): 261 - 290.
- [33] T. Tao, “Nonlinear Dispersive Equations,” Published for the Conference Board of the Mathematical Sciences, Washington, DC, 2006.
- [34] T. Tao and A. Vargas, *A bilinear approach to cone multipliers. I. Restriction estimates*, *Geom. Funct. Anal.*, **10** no. 1 (2000), 185–215.
- [35] T. Tao, M. Visan, and X. Zhang. "The nonlinear Schrödinger equation with combined power-type nonlinearities." *Comm. Partial Differential Equations*, **32** no. 7-9 (2007) :1281–1343.
- [36] T. Tao, M. Visan, and X. Zhang. "Minimal-mass blowup solutions of the mass-critical NLS." *Forum Mathematicum*, **20** no. 5 (2008) : 881 - 919.
- [37] T. Tao, M. Visan, and X. Zhang. "Global well-posedness and scattering for the defocusing mass - critical nonlinear Schrödinger equation for radial data in high dimensions." *Duke Mathematical Journal*, **140** no. 1 (2007) : 165 - 202.
- [38] M. E. Taylor, “Pseudodifferential Operators and Nonlinear PDE,” Birkhäuser, Boston, 1991.
- [39] M. E. Taylor, “Partial Differential Equations I - III,” Springer-Verlag, New York, 1996.
- [40] M. E. Taylor “Short time behavior of solutions to nonlinear Schrödinger equations in one and two space dimensions” *Comm. Partial Differential Equations* **31** (2006): 955 - 980.
- [41] M. E. Taylor, “Tools for PDE” American Mathematical Society, *Mathematical Surveys and Monographs* **31** Providence, RI, 2000.
- [42] M. Visan “The defocusing energy-critical nonlinear Schrödinger equation in higher dimensions” *Duke Mathematical Journal* **138** (2007): 281 - 374.
- [43] M. Weinstein, “Nonlinear Schrödinger equations and sharp interpolation estimates” *Communications in Mathematical Physics* **87** no. 4 (1982/83): 567 - 576.

- [44] M. Weinstein, "The nonlinear Schrödinger equation – singularity formation, stability and dispersion" *The connection between infinite - dimensional and finite - dimensional dynamical systems* (Boulder CO) **99** (1989): 213 - 232.
- [45] K. Yosida, "Functional Analysis" Grundlehren der Mathematischen Wissenschaften [Fundamental Principles of Mathematical Sciences], Volume 123, 6th Edition Springer - Verlag, Berlin, 1980.

UNIVERSITY OF CALIFORNIA, BERKELEY